

Two Permian lyttoniid brachiopods from Akasaka, central Japan

JUN-ICHI TAZAWA¹, TERUO ONO² and MASAKAZU HORI³

¹Department of Geology, Faculty of Science, Niigata University, Niigata 950-2181, Japan

²Honden 1552-141, Hozumi-cho, Motosu-gun, Gifu Prefecture 501-0236, Japan

³Hirui-cho 377-18, Ogaki, Gifu Prefecture 503-2216, Japan

Received 17 April 1998 ; Revised manuscript accepted 6 November 1998

Abstract. Two lyttoniid brachiopod species, *Coscinophora magnifica* Cooper and Grant, 1974 and *Leptodus nobilis* (Waagen, 1883) are described from the Middle and Upper Permian limestones in the Akasaka district, Mino Belt, central Japan. The Permian brachiopod fauna of Akasaka is characterized by a mixture of both Tethyan and North American elements. The fauna is considered to have occupied a mid-equatorial region of the Panthalassa Ocean in Permian time.

Key words : Akasaka, brachiopods, central Japan, *Coscinophora magnifica*, *Leptodus nobilis*, Mino Belt, Permian

Introduction

The Akasaka district, northwest of Ogaki City in the Gifu Prefecture is a famous, classical area for the Japanese Permian. The Permian marine invertebrate fossils of the Akasaka Limestone in this district have been studied by many authors since the pioneering works of Schwager (1883), Gottsche (1884) and Ozawa (1927). Brachiopods from the Akasaka Limestone were studied by Sato (1919), Ozawa (1927) and Hayasaka (1925, 1932), and ten species were described or listed: *Lyttonia richthofeni* Keyser (Sato, 1919; Hayasaka, 1925; Ozawa, 1927), *Enteleles akasakensis* Ozawa (Ozawa, 1927; Hayasaka, 1932), *Scacchinella* cf. *gigantea* Gemmellaro (Hayasaka, 1925, 1932), *Reticularia lineata* Martin, *R. waageni* Loczy, *R. cf. inaequilateralis* Gemmellaro, *Terebratuloidea*? sp. (Hayasaka, 1925), *Enteretes minoensis* Hayasaka, *E. suessi* Schellwien and *Geyerella* sp. (Hayasaka, 1932). Recently a lyttoniid species *Coscinophora magnifica* Cooper and Grant, from its lower part (*Parafusulina* Zone), was described and added to the list by Tazawa (1997b).

In this paper two lyttoniids from the Akasaka Limestone, *Coscinophora magnifica* Cooper and Grant, 1974 from its lower part (*Parafusulina* Zone) at Locality 1, Kinshozan, and *Leptodus nobilis* (Waagen, 1883) from its uppermost part (*Codonofusiella-Reichelina* Zone) at Locality 2, Hanaakayama, are described, and their palaeobiogeography is discussed. The fossil localities are shown on Figure 1. The first author (J.T.) is responsible for the systematic descriptions and discussion of Permian brachiopod palaeobiogeography, and the second and third authors (T.O. and M.H.) are respon-

sible for field geology and collecting the specimens. All the specimens described here are stored in the Department of Geology, Faculty of Science, Niigata University.

Distribution of *Coscinophora* and *Leptodus*

The distribution of the genus *Coscinophora* is restricted stratigraphically and geographically. It has been described from only two regions, the Lower Permian (Wolfcampian) and Middle Permian (Leonardian) of west Texas, U.S.A. and the Middle Permian (*Parafusulina* Zone) of Akasaka, central Japan. *Coscinophora magnifica* Cooper and Grant has been known from the Middle Permian (Road Canyon Formation) of west Texas and the Middle Permian (lower part of the Akasaka Limestone) of Akasaka, central Japan (Sato, 1919; Hayasaka, 1925, 1932; Cooper and Grant, 1974; Tazawa, 1997b, this paper).

On the other hand, the genus *Leptodus* is widely distributed in the Lower to Upper Permian of the Tethyan and its neighboring regions. *Leptodus nobilis* (Waagen) is known from the Middle and Upper Permian of Hungary (Schr ter, 1963), Croatia (Sremac, 1986), Serbia (Simic, 1933), Transcaucasia (Licharew, 1932; Sarytcheva, 1964; Ruzhentsev and Sarytcheva, 1965; Kotljar in Kotljar and Zakharov, 1989), Salt Range (Waagen, 1883; Noetling, 1904, 1905; Frech, 1911; Fredericks, 1916; Cooper and Grant, 1974) and Khisor Range (Grant, 1976) in Pakistan, Kumaon Himalayas (Diener, 1897), Kashmir (Diener, 1915), Cambodia (Mansuy, 1913, 1914; Termier and Termier, 1960; Chi-Thuan, 1961), Laos (Mansuy, 1912), Timor (Hamlet, 1928; Wanner and Sieverts, 1935), Port Keats in northern Australia (Thomas, 1957), Tibet (Zhan and



Figure 1. Map showing the fossil localities (using the topographical map of "Ogaki" scale 1 : 25,000 published by Geographical Survey of Japan).

Wu, 1982), Yunnan (Fang and Fan, 1994), Sichuan (Hayasaka, 1917, 1922b; Huang, 1932), Guizhou (Hayasaka, 1917, 1922b; Huang, 1932; Feng and Jiang, 1978), Guangxi (Huang, 1936; Yang *et al.*, 1977), Guangdong (Yang *et al.*, 1977; Zhan, 1979), Hunan (Yang *et al.*, 1977; Liao and Meng, 1986) and Hubei (Yang *et al.*, 1977; Yang, 1984) in South China, Jiangxi (Hayasaka, 1922b), Fujian (Wang *et al.*, 1982; Zhu, 1990) and Zhejiang (Liang, 1990) in East China, Gansu (Zhang *et al.*, 1983) and Qinghai (Jin *et al.*, 1979) in Northwest China, Inner Mongolia (Grabau, 1931; Lee and Gu, 1976; Lee *et al.*, 1980; Duan and Li, 1985; Gu and Zhu, 1985), Heilongjiang (Lee *et al.*, 1980) and Jilin (Lee *et al.*, 1980) in Northeast China, South Primorye in eastern Russia (Licharew and Kotljar, 1978), South Kitakami Belt (Yabe, 1900; Hayasaka, 1917, 1922a; Tazawa, 1976, 1987; Minato *et al.*, 1979), Hida Gaien Belt (Tazawa, 1987; Tazawa and Matsumoto, 1998), Maizuru Belt (Mashiko, 1934; Shimizu, 1961) and Mino Belt (this paper) in Japan.

Permian reconstruction of the Akasaka Limestone

From the above data, it is clear that the genus *Leptodus* is a Tethyan element and the genus *Coscinophora* is a North American element, and the Permian brachiopod fauna of the Akasaka Limestone is a mix of Tethyan and North American elements.

The Akasaka Limestone is one of the limestone-greenstone blocks in the Jurassic melange of the Mino Belt in central Japan. The limestone-greenstone blocks are con-

sidered to have originated from reef-seamount complexes formed at the equatorial region of the Panthalassa Ocean in Permian time on the basis of palaeomagnetic (Hattori and Hirooka, 1977, 1979), sedimentological (Sano, 1988; Sano and Kanmera, 1996) and palaeontological data (Ishii *et al.*, 1985; Ozawa, 1987; Tazawa, 1991, 1992). However, none of these authors gave their opinion about the palaeolatitude of the Permian reef-sea mount complexes.

Tazawa (1997a, b) and Tazawa and Shen (1997) previously mentioned that these limestone-greenstone blocks were formed at the mid-equatorial region of Permian Panthalassa based on palaeobiogeographical studies of brachiopod faunas of the Hiyomo and Akasaka districts, Mino Belt. The occurrence of a typical Tethyan-type genus *Leptodus* together with a North American-type genus *Coscinophora* from the Akasaka Limestone strongly supports the above opinion.

Systematic descriptions

- Order Productida Waagen, 1883
- Suborder Strophalosiidina Waagen, 1883
- Superfamily Lytonioidea Waagen, 1883
- Family Lytoniidae Waagen, 1883
- Subfamily Lytoniinae Waagen, 1883
- Genus *Coscinophora* Cooper and Stehli, 1955

Type species.—*Coscinophora nodosa* Cooper and Stehli, 1955

Coscinophora magnifica Cooper and Grant, 1974

Figures 2-3-5, 3

Coscinophora magnifica Cooper and Grant, 1974, p. 454, pl. 182, figs. 35, 36; pl. 184, figs. 1-8; pl. 185, figs. 1-17; pl. 186, figs. 1-13; pl. 187, figs. 1-11; pl. 188, figs. 15-22, 24; Tazawa, 1997b, p. 447, figs. 2-1-3.

Lyttonia richthofeni (Kayser). Sato, 1919, p. 276, text-fig.; Hayasaka, 1932, p. 1.

Lyttonia sp. Hayasaka, 1925, p. 143, text-fig.

Material.—Three specimens, NU-B110-112, from the lower part (*Parafusulina* Zone) of the Akasaka Limestone at Locality 1, Kinshozan, Akasaka.

Description.—Shell large for genus, transversely subelliptical in outline; length 56 mm, width 65 mm in the largest specimen (NU-B111). Pedicle valve slightly convex in lateral and anterior profiles except anterolateral margins which form steep slopes.

Pedicle valve interior with a median ridge and at least 12 lateral ridges, both of them consisting of a row of small beads. Lateral ridges rather regularly and symmetrically arranged on both sides of median ridge, nearly straight but slightly arched facing convexity towards front. Number and size of beads are obscure in the present material.

Brachial valve interior partly preserved in one specimen (NU-B110). Lateral lobes rather regularly and symmetrically arranged anteriorly on both sides of median lobe, but irregu-

larly developed posteriorly. Each lateral lobe with a row of small, subrectangular holes, numbering 4 holes per 10 mm (Figure 3). Other internal structures not observed.

Remarks.—The present specimens are assigned to the genus *Coscinophora* on the basis of their pedicle valve moniliseptate, median and lateral ridges consisting of a row of beads, and their brachial valve median and lateral lobes consisting of a row of holes. These specimens are referred to *Coscinophora magnifica* Cooper and Grant, 1974, from the Road Canyon Formation of the Glass Mountains, west Texas, on account of the comparable size and shape of shell, and size and number of holes on the lateral lobes of the brachial valve.

The lyttoniid brachiopods collected from the lower part (Kasumi Zone of Wakimizu, 1902) of the Akasaka Limestone and reported as *Lyttonia richthofeni* (Kayser, 1883) (Sato, 1919, p. 276; Hayasaka, 1932, p. 1) and *Lyttonia* sp. (Hayasaka, 1925, p. 143) may be conspecific with the present species.

Genus *Leptodus* Kayser, 1883

Type species.—*Leptodus richthofeni* Kayser, 1883

Leptodus nobilis (Waagen, 1883)

Figures 2-1, 2-2, 4

Lyttonia nobilis Waagen, 1883, p. 398, pl. 29, figs. 1-3; pl. 30, figs. 1, 2, 5, 6, 8, 10, 11; Diener, 1897, p. 37, pl. 1, figs. 5-7;




Figure 2. 1, 2. *Leptodus nobilis* (Waagen, 1883). Internal mould of a pedicle valve and the latex cast, NU-B128. 3-5. *Coscinophora magnifica* Cooper and Grant, 1974. Internal moulds of three pedicle valves, 3: NU-B110; 4: NU-B111; 5: NU-B112. (All figures are natural size)



Figure 3. A part of the brachial valve of *Coscinophora magnifica* Cooper and Grant, 1974, showing details of the lateral lobes, NU-B110. ($\times 3$)

Noetling, 1904, p. 112, text-figs. 4-7; Noetling, 1905, p. 140, pl. 17, figs. 1, 2; pl. 18, figs. 1-11, text-fig. 2; Mansuy, 1913, p. 123, pl. 13, fig. 10; Mansuy, 1914, p. 32, pl. 6, figs. 7a-d; pl. 7, figs. 1a-e; Diener, 1915, p. 99, pl. 10, fig. 15; Grabau, 1931, p. 285, pl. 28, figs. 4, 5 only; Huang, 1932, p. 89, pl. 7, figs. 9, 10; pl. 8, figs. 8, 9; pl. 9, figs. 1-8, text-figs. 8-11; Simic, 1933, p. 49, pl. 4, fig. 1.

Lyttonia tenuis Waagen, 1883, p. 401, pl. 30, figs. 3, 4, 7, 9.

Lyttonia sp. Yabe, 1900, p. 2, text-figs. 1, 2.

Lyttonia cf. *tenuis* Waagen. Mansuy, 1912, p. 19, pl. 4, fig. 4; pl. 5, figs. 1a-e; Huang, 1936, p. 493, pl. 1, fig. 6.

Oldhamina (*Lyttonia*) *richthofeni* var. *nobilis* Waagen. Fredericks, 1916, p. 76, pl. 4, fig. 2, text-fig. 22.

Lyttonia richthofeni (Kayser). Frech, 1911, p. 135, pl. 20, figs. 2a, b only; Hayasaka, 1917, p. 43, pl. 18, figs. 1-8; Hayasaka, 1922a, p. 62, pl. 11, figs. 1-6; Hayasaka, 1922b, p. 103, pl. 4, figs. 12, 13; Licharew, 1932, p. 56, 86, pl. 1, figs. 1-16; pl. 2, figs. 1, 2, 5, 7, 10, 12; pl. 3, figs. 2-7; pl. 4, figs. 1-17; pl. 5, figs. 1-4, 6; Mashiko, 1934, p. 182, text-fig.

Lyttonia (*Leptodus*) *richthofeni* Kayser. Hamlet, 1928, p. 31, pl. 6, figs. 1-4.

Lyttonia richthofeni forma *nobilis* Waagen. Licharew, 1932, p. 69, 96, pl. 2, figs. 13, 14; pl. 5, figs. 1-4, 6, text-fig. 3.

Leptodus nobilis (Waagen). Wanner and Sieverts, 1935, p. 249, pl. 9, figs. 27, 28, text-figs. 16-18; Termier and Termier, 1960, p. 241, text-pl. 3, figs. 1-10; Chi-Thuan, 1961, p. 274, pl. 1, figs. 1a, b; Schr ter, 1963, p. 107, pl. 3, figs. 5-8; Sarytcheva, 1964, p. 65, pl. 7, figs. 5-8, text-fig. 1; Ruzhentsev and Sarytcheva, 1965, pl. 39, figs. 6-8; Cooper and Grant, 1974, pl. 191, figs. 8, 9; Grant, 1976, pl. 43, figs. 18, 19; Lee and Gu, 1976, p. 267, pl. 162, figs. 1, 2; Tazawa, 1976, pl. 2, fig. 8; Yang *et al.*, 1977, p. 371, pl. 147, fig. 5; Feng and Jiang, 1978, p. 269, pl. 100, fig. 2; Licharew and Kotljar, 1978, pl. 14, figs. 13-15; Jin *et al.*, 1979, p. 82, pl. 23, fig. 15; Minato *et al.*, 1979, pl. 66, figs. 1, 4, 5; Zhan, 1979, p. 93, pl. 9, fig. 12; Lee *et al.*, 1980, p. 389, pl. 172, figs. 15, 16; Wang *et al.*, 1982, p. 229, pl. 95, fig. 20; Zhan and Wu, 1982, pl. 4, fig. 4; Zhang *et al.*, 1983, p. 297, pl. 102, figs. 7, 8; Yang, 1984, p. 226, pl. 35, fig. 12; Gu and Zhu, 1985, pl. 1, figs. 31,

33, 34; Liao and Meng, 1986, p. 81, pl. 2, figs. 24, 25; Sremac, 1986, p. 30, pl. 10, figs. 1-2; Tazawa, 1987, fig. 1-11; Kotljar in Kotljar and Zakharov, 1989, pl. 20, fig. 6; pl. 23, fig. 12; Liang, 1990, p. 225, pl. 40, figs. 1, 5; Fang and Fan, 1994, p. 83, pl. 23, figs. 1-3; pl. 30, fig. 5; Tazawa and Matsumoto, 1998, p. 7, pl. 2, figs. 7-12.

Lyttonia cf. *nobilis* Waagen. Huang, 1936, p. 493, pl. 1, fig. 5.

Leptodus cf. *nobilis* (Waagen). Thomas, 1957, p. 177, pl. 20, figs. 1-6.

Leptodus richthofeni Kayser. Shimizu, 1961, pl. 18, figs. 14, 15; Schr ter, 1963, p. 106, pl. 3, fig. 4; Sarytcheva, 1964, p. 65, pl. 7, figs. 2-4; Yang *et al.*, 1977, p. 372, pl. 147, fig. 10; Yang, 1984, p. 226, pl. 35, fig. 11; Duan and Li, 1985, p. 119, pl. 35, figs. 17-19.

Leptodus ivanovi Fredericks. Minato *et al.*, 1979, pl. 66, fig. 3.

Leptodus sp. Minato *et al.*, 1979, pl. 66, fig. 2.

Leptodus tenuis (Waagen). Duan and Li, 1985, p. 119, pl. 35, figs. 14-16; Liang, 1990, p. 226, pl. 40, fig. 9; Zhu, 1990, p. 79, pl. 18, figs. 19-21; Fang and Fan, 1994, p. 83, pl. 23, figs. 4-5; pl. 30, fig. 6.

Leptodus sp. Tazawa, 1987, fig. 1-10.

Material.—One specimen, NU-B128, from the uppermost part (*Codonofusiella-Reichelina* Zone) of the Akasaka Limestone at Locality 2, Hanaokayama, Akasaka.

Description.—Shell medium to small for genus, transversely subtrigonal in outline, with flattened pedicle valve, having greatest width near anterior margin; length 31 mm, width 53 mm.

Pedicle valve interior with regularly and symmetrically arranged lateral septa on both sides of median septum. Median septum extended for whole length of pedicle valve with blunt, rounded edge. Lateral septa rounded, solid (Figure 4), nearly straight, but slightly arched toward front, numbering 11 in the pedicle valve. Other internal structures not observed.

Remarks.—This specimen is safely assigned to the genus *Leptodus* by its rounded and solid (solidiseptate) lateral septa in the pedicle valve. The Akasaka specimen may be an immature one of *Leptodus nobilis* (Waagen), originally described and figured by Waagen (1883, p. 398, pl. 29, figs. 1-3; pl. 30, figs. 1, 2, 5, 6, 8, 10, 11) from the Wargal and Chhidru Formations of the Salt Range, Pakistan.

The single lyttoniid specimen figured by Ozawa (1927, pl. 45, fig. 12) as *Lyttonia richthofeni* Kayser from the lower part (Same Zone of Wakimizu, 1902) of the Akasaka Limestone is clearly distinguished from the present species by its grooved (angustilobate) lateral septa in the pedicle valve, which is the diagnostic character of the genus *Eolyttonia* Fredericks,



Figure 4. Longitudinal profile of the pedicle valve of *Leptodus nobilis* (Waagen, 1883). Right lateral view of a section of the latex cast, showing the lateral septa, NU-B128. ($\times 2$)

1924.

Acknowledgments

We thank Neil W. Archbold of Deakin University, Clayton, Australia for critical reading of the manuscript; Shen Shu-zhong of the same university for photography.

References cited

- Chi-Thuan, T.T., 1961: Les brachiopodes permians du Phnom-Tup (Sisophon-Cambodge). *Annales de la Faculté des Sciences, Université de Saigon*, 1961, p. 267-308.
- Cooper, G.A. and Grant, R.E., 1974: Permian brachiopods of West Texas, 2. *Smithsonian Contributions to Paleobiology*, no. 15, p. 233-793.
- Cooper, G.A. and Stehli, F.G., 1955: New genera of Permian brachiopods from West Texas. *Journal of Paleontology*, vol. 29, no. 3, p. 469-474.
- Diener, C., 1897: Himalayan fossils; The Permocarboniferous fauna of Chitichun, No. 1. *Palaeontologia Indica, Series 15*, vol. 1, pt. 3, p. 1-104.
- Diener, C., 1915: The Anthracolithic faunas of Kashmir, Kanaur and Spiti. *Palaeontologia Indica, New Series*, vol. 5, no. 2, p. 1-135.
- Duan, C. and Li, W., 1985: Description of fossils, (4) Phylum Brachiopoda. In, Ding, Y., Xia, G., Duan, C., Li, W., Liu, X. and Liang, Z., Study on the Early Permian stratigraphy and fauna in Zhesi district, Nei Mongol Zizhiqu (Inner Mongolia). *Bulletin of the Tianjin Institute of Geology and Mineral Resources*, no. 10, p. 99-145, 199-214. (in Chinese)
- Fang, R. and Fan, J., 1994: Middle to Upper Carboniferous-Early Permian Gondwana facies and palaeontology in western Yunnan, 121 p. Yunnan Science and Technology Press, Kunming. (in Chinese)
- Feng, R. and Jiang, Z., 1978: Phylum Brachiopoda. In, Geological and Palaeontological Team of Guizhou ed., *Palaeontological atlas of Southwest China; Guizhou, pt. 2. Carboniferous to Quaternary volume*, p. 231-305, Geological Publishing House, Beijing. (in Chinese)
- Frech, F., 1911: Die Dyas. In, Richthofen, F. von ed., *China, Fünfter Band*, p. 103-137, Dietrich Reimer, Berlin.
- Fredericks, G., 1916: The palaeontological notes, 2. On some Upper Palaeozoic Brachiopoda of Eurasia. *Trudy Geologicheskogo Komiteta, Novaya Seriya*, fascicle 156, p. 1-87. (in Russian)
- Fredericks, G., 1924: Upper Paleozoic of the Ussuriland, 1. Brachiopoda. *Materialy po geologii i poleznym iskopaemym Dalnego Vostoka*, no. 28, p. 1-52. (in Russian)
- Gottsche, C., 1884: Ueber japanisches Carbon. *Zeitschrift der Deutschen Geologischen Gesellschaft*, vol. 36, p. 653-654.
- Grabau, A.W., 1931: The Permian of Mongolia. In, Reeds, C.A., ed., *Natural history of Central Asia*, vol. 4, 665 p. American Museum of Natural History, New York.
- Grant, R.E., 1976: Permian brachiopods from southern Thailand. *The Paleontological Society, Memoir*, 9, p. 1-269.
- Gu, F. and Zhu, R., 1985: Lower Permian brachiopods from Lin-Dong, Nei Mongol. *Bulletin of the Shenyang Institute of Geology and Mineral Resources*, no. 12, p. 74-97. (in Chinese)
- Hamlet, B., 1928: Permische Brachiopoden, Lamellibranchiaten und Gastropoden von Timor. *Jaarboek van het Mijneuzen in Nederlandsch-Indie*, vol. 56, no. 2, p. 1-115.
- Hattori, I. and Hirooka, K., 1977: Paleomagnetic study of greenstone in the Mugi-Kamiaso area, Gifu Prefecture, central Japan. *Journal of the Japanese Association of Mineralogists, Petrologists and Economic Geologists*, vol. 72, no. 9, p. 340-353.
- Hattori, I. and Hirooka, K., 1979: Paleomagnetic results from Permian greenstones in central Japan and their geologic significance. *Tectonophysics*, vol. 57, p. 211-235.
- Hayasaka, I., 1917: On the brachiopod genus *Lyttonia* with several Japanese and Chinese examples. *Journal of the Geological Society of Tokyo*, vol. 24, no. 288, p. 43-53.
- Hayasaka, I., 1922a: Some Permian brachiopods from the Kitakami Mountains. *Japanese Journal of Geology and Geography*, vol. 1, no. 2, p. 51-70.
- Hayasaka, I., 1922b: Paleozoic Brachiopoda from Japan, Korea and China, Part 1. Middle and Southern China. *Science Reports of the Tohoku Imperial University, Second Series*, vol. 6, no. 1, p. 1-116.
- Hayasaka, I., 1925: On *Lyttonia* and several other brachiopods from Kinsho-zan. *Journal of the Geological Society of Tokyo*, vol. 32, no. 379, p. 142-146. (in Japanese)
- Hayasaka, I., 1932: On three brachiopod species of the subfamily Orthotetinae in the *Fusulina*-Limestone of Kinshozan, Akasaka-Mati, Prov. Mino, Japan. *Memoir of the Faculty of Science and Agriculture, Taihoku Imperial University, Formosa, Japan*, vol. 4, nos. 2/3, p. 1-7.
- Huang, T.K., 1932: Late Permian Brachiopoda of southwestern China. *Palaeontologia Sinica, Series B*, vol. 9, fascicle 1, p. 1-139.
- Huang, T.K., 1936: On the occurrence of *Lyttonia* in the Wolfcamp Series of the Glass Mountains of Texas with notes on lyttonids from southwestern China. *Bulletin of the Geological Society of China*, vol. 15, no. 1, p. 489-493.
- Ishii, K., Okimura, Y. and Ichikawa, K., 1985: Notes on Tethys biogeography with reference to Middle Permian fusulinaceans. In, Nakazawa, K. and Dickins, J.M. eds., *The Tethys: Her paleogeography and paleobiogeography from Paleozoic to Mesozoic*, p. 139-155, Tokai University Press, Tokyo.
- Jin, Y., Ye, S., Yu, H. and Sun, D., 1979: Phylum Brachiopoda. In, Nanjing Institute of Geology and Palaeontology and Qinghai Institute of Geological Sciences eds., *Palaeontological Atlas of Northwest China; Qinghai, pt. 1*, p. 60-217, Geological Publishing House, Beijing. (in Chinese)
- Kayser, E., 1883: Obercarbonische Fauna von Lo-Ping. In, Richthofen, F. von ed., *China, Vierter Band*, p. 160-208, Dietrich Reimer, Berlin.
- Kotljars, G.V. and Zakharov, Yu. D., 1989: *Evolution of the latest Permian biota; Midian regional stage in the USSR*, 182 p. Nauka, Leningrad. (in Russian)
- Lee, L. and Gu, F., 1976: Carboniferous and Permian Bra-

- chiopoda. In, Geological Bureau of Nei Mongol and Geological Institute of Northeast China eds., *Palaeontological Atlas of Northeast China; Nei Mongol, pt. 1. Palaeozoic volume*, p. 228-306, Geological Publishing House, Beijing. (in Chinese)
- Lee, L., Gu, F. and Su, Y., 1980: Carboniferous and Permian Brachiopoda. In, Shenyang Institute of Geology and Mineral Resources ed., *Palaeontological Atlas of Northeast China, pt. 1. Palaeozoic volume*, p. 327-428, Geological Publishing House, Beijing. (in Chinese)
- Liang, W., 1990: Lengwu Formation of Permian and its brachiopod fauna in Zhejiang Province. *Geological Memoirs, Ministry of Geology and Mineral Resources, People's Republic of China, Series 2*, no. 10, p. 1-522. (in Chinese)
- Liao, Z. and Meng, F., 1986: Late Chanxianian brachiopods from Huatang of Chen Xian County, southern Hunan. *Memoirs of Nanjing Institute of Geology and Palaeontology*, no. 22, p. 71-94. (in Chinese)
- Licharew, B.K., 1932: Fauna of the Permian deposits of Northern Caucasus, 2. Brachiopoda; Family Lyttonidae Waagen. *Trudy Vsesoyuznogo Geologo-Razvedochnogo Obedicheniya NKTP SSSR*, fascicle 215, p. 55-111. (in Russian)
- Licharew, B.K. and Kotjar, G.V., 1978: Permskie brachiopody Yuzhnogo Primorya. In, Popeko, L.I., ed., *Verkhniy Paleozoy Severo-Vostochnoy Azii*, p. 63-75, Dalnevostochniy Nauchnyy Tsentr, Akademiya Nauk SSSR, Vladivostok. (in Russian)
- Mansuy, H., 1912: Mission du Laos. *Mémoires du Service Géologique de L'Indochine*, vol. 1, fascicle 4, p. 1-52.
- Mansuy, H., 1913: Faunes des Calcaires à Productus de L'Indochine, Première Série. *Mémoires du Service Géologique de L'Indochine*, vol. 2, fascicle 4, p. 1-133.
- Mansuy, H., 1914: Faunes des Calcaires à Productus de L'Indochine, Deuxième Série. *Mémoires du Service Géologique de L'Indochine*, vol. 3, fascicle 3, p. 1-59.
- Mashiko, K., 1934: Discovery of *Lyttonia* in a limestone exposed at Takauti, Nakayakuno-mura, Amata-gun, Kyoto Prefecture. *Japanese Journal of Geology and Geography*, vol. 11, nos. 3/4, p. 181-183.
- Minato, M., Hunahashi, M., Watanabe, J. and Kato, M. eds., 1979: *Variscan geohistory of Northern Japan: The Abean Orogeny*, 427 p. Tokai University Press, Tokyo.
- Noetling, F., 1904: Über den Bau und die Organisation der Lyttoniidae Waagen. *Verhandlungen der Deutschen Zoologischen Gesellschaft*, 1904, p. 103-122.
- Noetling, F., 1905: Untersuchungen über die Familie Lyttoniidae Waagen. emend. Noetling. *Palaeontographica*, vol. 51, p. 129-154.
- Ozawa, T., 1987: Permian fusulinacean biogeographic provinces in Asia and their tectonic implications. In, Taira, A. and Tashiro, M., eds.: *Historical biogeography and plate tectonic evolution of Japan and eastern Asia*, p. 45-63, Terra Scientific Publishing Company, Tokyo.
- Ozawa, Y., 1927: Stratigraphical studies of the *Fusulina* Limestone of Akasaka, Province of Mino. *Journal of the Faculty of Science, Imperial University of Tokyo, Section 2*, vol. 2, pt. 3, p. 121-164.
- Ruzhentsev, V.E. and Sarytcheva, T.G., 1965: Razvitie i smena morskikh organizmov na rubezhe Paleozoya i Mezozoya. *Trudy Paleontologicheskogo Instituta*, vol. 108, p. 1-431. (in Russian)
- Sano, H., 1988: Permian oceanic-rocks of Mino Terrane, central Japan, part 2. Limestone facies. *Journal of the Geological Society of Japan*, vol. 94, no. 12, p. 963-976.
- Sano, H. and Kanmera, K., 1996: Microbial controls on Panthalassan Carboniferous-Permian oceanic buildups, Japan. *Facies*, vol. 34, p. 239-256.
- Sarytcheva, T.G., 1964: Oldhaminid brachiopods from the Permian of Transcaucasia. *Paleontologicheskii Zhurnal*, 1964, no. 3, p. 58-72. (in Russian)
- Sato, D., 1919: On *Lyttonia* recently discovered at Akasaka, Province Mino. *Journal of the Geological Society of Tokyo*, vol. 26, no. 309, p. 276-277. (in Japanese)
- Schréter, Z., 1963: A Bükkhegység felső-permi brachiopodai. *Geologica Hungarica, Series Palaeontologica*, fascicle 28, p. 1-181.
- Schwager, C., 1883: Carbonische Foraminiferen aus China und Japan. In, Richthofen, F. von ed., *China, Vierter Band*, p. 106-159, Dietrich Reimer, Berlin.
- Shimizu, D., 1961: Brachiopod fossils from the Permian Maizuru Group. *Memoirs of the College of Science, University of Kyoto, Series B*, vol. 27, no. 3, p. 309-350.
- Simic, V., 1933: Gornji Perm u Zapadnoj Srbiji. *Mémoires du Service Géologique du Royaume de Yougoslavie*, vol. 1, p. 1-130.
- Sremac, J., 1986: Middle Permian brachiopods from the Velebit Mts. (Croatia, Yugoslavia). *Palaeontologia Jugoslavica*, vol. 35, p. 1-43.
- Tazawa, J., 1976: The Permian of Kesennuma, Kitakami Mountains: A preliminary report. *Earth Science (Chikyū Kagaku)*, vol. 30, no. 3, p. 175-185.
- Tazawa, J., 1987: Permian brachiopod faunas of Japan and their palaeobiogeography. *Gekkan Chikyū*, vol. 9, no. 5, p. 252-255. (in Japanese)
- Tazawa, J., 1991: Middle Permian brachiopod biogeography of Japan and adjacent regions in East Asia. In, Ishii, K., Liu, X., Ichikawa, K. and Huang, B. eds., *Pre-Jurassic geology of Inner Mongolia, China: Reports of China-Japan Cooperative Research Group, 1987-1989*, p. 213-230, Matsuya Insatsu, Osaka.
- Tazawa, J., 1992: Middle Permian brachiopod faunas in East Asia and their zoogeographic significance. *Journal of the Geological Society of Japan*, vol. 98, no. 6, p. 483-496. (in Japanese)
- Tazawa, J., 1997a: Permian brachiopods from Hiyomo, east of Takayama, central Japan: A palaeobiogeographical data on the origin of limestone-greenstone blocks in the Mino Belt. *Journal of the Geological Society of Japan*, vol. 103, no. 9, p. 908-911. (in Japanese)
- Tazawa, J., 1997b: *Coscinophora* (Permian Brachiopoda) from the Akasaka Limestone, Mino-Belt, central Japan and its palaeobiogeographical significance. *Earth Science (Chikyū Kagaku)*, vol. 51, no. 6, p. 447-451. (in Japanese)
- Tazawa, J. and Matsumoto, T., 1998: Middle Permian brachiopods from the Oguradani Formation, Ise district, Hida Gaian Belt, central Japan. *Science Reports of Niigata University, Series E*, no. 13, p. 1-19.
- Tazawa, J. and Shen, S., 1997: Middle Permian brachiopods from Hiyomo, Mino Belt, central Japan: Their provincial relationships with North America. *Science Reports of Niigata University, Series E*, no. 12, p. 1-17.
- Termier, H. and Termier, G., 1960: Contribution à la classifi-

- cation des Brachiopodes: le lophophore des Collolophides nov. ord.; Appendice. Les Oldhaminidés du Cambodge. *Société Géologique de France, Bulletin, Série 7*, vol. 1, no. 3, p. 233-243.
- Thomas, G.A., 1957: Oldhaminid brachiopods in the Permian of Northern Australia. *Journal of the Palaeontological Society of India*, vol. 2, p. 174-182.
- Waagen, W., 1883: Salt Range fossils; *Productus*-Limestone fossils. *Palaeontologia Indica, Series 13*, vol. 1, pt. 4, p. 391-546.
- Wakimizu, T., 1902: On the Akasaka Limestone, Prov. Mino. *Journal of the Geological Society of Tokyo*, vol. 9, no. 102, p. 71-75; no. 104, p. 163-169; no. 105, p. 205-212; no. 108, p. 231-239. (*in Japanese*)
- Wang, G., Liu, Q., Jin, Y., Hu, S., Liang, W. and Liao, Z., 1982: Phylum Brachiopoda. In, Nanjing Institute of Geology and Mineral Resources ed., *Palaeontological Atlas of East China, pt. 2. Late Palaeozoic volume*, p. 186-256, Geological Publishing House, Beijing. (*in Chinese*)
- Wanner, J. and Sieverts, H., 1935: Zur Kenntnis der permischen Brachiopoden von Timor, 1. Lyttoniidae und ihre biologische und stammesgeschichtliche Bedeutung. *Neues Jahrbuch für Mineralogie, Geologie und Paläontologie*, vol. 74, p. 201-281.
- Yabe, H., 1900: The brachiopod *Lyttonia* from Rikuzen Province. *Journal of the Geological Society of Tokyo*, vol. 7, no. 79, p. 1-4.
- Yang, D., 1984: Systematic description of Palaeontology: Brachiopoda. In, Yichang Institute of Geology and Mineral Resources ed., *Biostratigraphy of the Yangtze Gorge area*, vol. 3, p. 203-239, 330-333, Geological Publishing House, Beijing. (*in Chinese*)
- Yang, D., Ni, S., Chang, M. and Zhao, R., 1977: Phylum Brachiopoda. In, Geological Institute of Hubei et al., eds., *Palaeontological atlas of South-central China, part 2. Late Palaeozoic volume*, p. 303-470, Geological Publishing House, Beijing. (*in Chinese*)
- Zhan, L., 1979: Description of fossils, (2) Brachiopoda. In, Hou, H., Zhan, L., Chen, B. and others, *The coal-bearing strata and fossils of Late Permian from Guangtung*, p. 61-100, Geological Publishing House, Beijing. (*in Chinese*)
- Zhan, L. and Wu, R., 1982: Early Permian brachiopods from Xainza district, Xizang (Tibet). In, CGQXP Editorial Committee, Ministry of Geology and Mineral Resources, ed., *Contribution to the geology of the Qinghai-Xizang (Tibet) Plateau*, vol. 7, p. 86-109, Geological Publishing House, Beijing. (*in Chinese*)
- Zhang, Y., Fu, L., Ding, P. and Qi, W., 1983: Phylum Brachiopoda. In, Xian Institute of Geology and Mineral Resources ed., *Palaeontological Atlas of Northwest China; Shaanxi, Gansu and Ningxia volume, pt. 2. Upper Palaeozoic*, p. 244-425, Geological Publishing House, Beijing. (*in Chinese*)
- Zhu, T., 1990: *The Permian coal-bearing strata and palaeobiocoenosis of Fujian*, 127 p. Geological Publishing House, Beijing. (*in Chinese*)